


50-year-old puzzle about β -decay rates resolved from first principles

Objectives

- Address the long-standing puzzle of why computations of β -decay rates in atomic nuclei are faster than what's expected from the β -decay of the free neutron
- Utilize state-of-the-art interactions from chiral effective-field-theory and computational methods to address the puzzle
- Explore the role of the coupling of the weak force to two nucleons and of strong correlations in the nucleus

Impact

- β -decay is the dominant decay mode of atomic nuclei
- β -decay rates enter models of heavy element synthesis in neutron star mergers and supernovae explosions
- Understanding β -decay relevant for neutrino-less double- β -decay, to reduce the uncertainty in extracting the neutrino mass scale



Accomplishments

- Resolved the long-standing discrepancy between experimental and theoretical β -decay rates from first principles.
- The coupling of the weak force to two nucleons and a proper treatment of strong correlations in the nucleus are necessary to correctly describe β -decay rates from light nuclei to the heavy nucleus ^{100}Sn

Caption: Gamow–Teller strength in ^{100}Sn . Comparison of the Gamow–Teller strength $|M_{\text{GT}}|^2$ for the β -decay of ^{100}Sn calculated in this work compared to experiment (Expt), and other models. Open symbols represent results obtained with the standard Gamow–Teller operator, filled symbols include two-body currents (2BCs) and partially filled symbols show values following from the multiplication of the computed Gamow–Teller strength by a phenomenological quenching factor.